



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2008

---

## **Coronary artery ectasia causing ischemia**

Husmann, L ; Herzog, B A ; Burkhard, N ; Valenta, I ; Weber, K ; Kaufmann, P A

DOI: <https://doi.org/10.1007/s00259-008-0895-x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-14227>

Journal Article

Published Version

Originally published at:

Husmann, L; Herzog, B A; Burkhard, N; Valenta, I; Weber, K; Kaufmann, P A (2008). Coronary artery ectasia causing ischemia. *European Journal of Nuclear Medicine and Molecular Imaging*, 35(11):2142.

DOI: <https://doi.org/10.1007/s00259-008-0895-x>

## Coronary artery ectasia causing ischemia


Lars Husmann · Bernhard A. Herzog · Nina Burkhard ·  
Ines Valenta · Klaus Weber · Philipp A. Kaufmann

Received: 19 June 2008 / Accepted: 29 June 2008 / Published online: 6 August 2008  
© Springer-Verlag 2008

A 46-year-old male smoker (70 kg, 175 cm) with no other cardiovascular risk factors presented with recurrent episodes of chest discomfort and breathing distress at physical exercise. Cardiac stress testing on a treadmill ergometer was clinically pathologic at 100 W, and the patient was subsequently referred to non-invasive hybrid cardiac imaging.

<sup>99m</sup>Tc-Tetrofosmin single-photon emission computed tomography (SPECT) images were acquired using a 1-day adenosine-stress/rest protocol and fused with a low-dose, prospectively gated computed tomography coronary angiography (effective radiation dose 1.6 mSv) [1, 2]. The latter revealed ectasia in the left anterior descending, the circumflex, and the right coronary artery (LAD, CX, RCA in panel B), but no relevant coronary artery stenosis. However, a reversible perfusion defect (ischemia) in the anterior myocardium was demonstrated by adenosine-stress SPECT (bluish colors in panel A; rest scan not shown).

Both ectasia and the absence of significant coronary artery stenosis were confirmed by invasive coronary angiography, suggesting conservative treatment [3].



L. Husmann · B. A. Herzog · N. Burkhard · I. Valenta ·  
K. Weber · P. A. Kaufmann (✉)  
Cardiovascular Center, University Hospital Zurich,  
Raemistrasse 100,  
8091 Zurich, Switzerland  
e-mail: pak@usz.ch

P. A. Kaufmann  
Zurich Center for Integrative Human Physiology,  
University of Zurich,  
Zurich, Switzerland

### References

1. Gaemperli O, Schepis T, Valenta I, et al. Cardiac image fusion from stand-alone SPECT and CT: clinical experience. *J Nucl Med*. 2007;48:696–703.
2. Husmann L, Valenta I, Gaemperli O, et al. Feasibility of low-dose coronary CT angiography: first experience with prospective ECG-gating. *Eur Heart J* 2008;29:191–7.
3. Kruger D, Stierle U, Herrmann G, et al. Exercise-induced myocardial ischemia in isolated coronary artery ectasias and aneurysms (“dilated coronopathy”). *J Am Coll Cardiol* 1999;34:1461–70.